

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

d88ac99f9883a0c44138fe74cb5af9a7e932d7255c1fc698308ed1ac5a5f92db

To view the reconstructed contents, please SCROLL DOWN to next page.

Seminar

Institute for Plasma Research

Title : Sculpted Ultracold Neutral Plasmas

Speaker: Dr. Vikram Dharodi

Michigan State University, USA

Date : 23rd October 2020 (Friday)

Time : 3.30 PM

Venue : Online - Join the talk:

https://meet.ipr.res.in/Dr.VikramDharodiv_PDFtalk

Abstract :

Ultracold neutral plasma (UNP) experiments allow for careful control of plasma properties across Coulomb coupling regimes. Here, we examine how UNPs can be used to study heterogeneous, non-equilibrium phenomena, including non-linear waves, transport, hydrodynamics, kinetics, stopping power and instabilities. Through a series of molecular dynamics simulations, we have explored UNPs formed with spatially modulated ionizing radiation. We have developed a computational model for such sculpted ultracold neutral plasmas that includes an ionic screened Coulomb interaction with a spatiotemporal screening length, and Langevin-based spatial ion-electron and ion-neutral collisions. We have also developed a hydrodynamics model and have extracted its field quantities (density, flow velocity and temperature) from the molecular dynamics simulation data, allowing us to investigate kinetics by examining moment ratios and phase-space dynamics; we find that it is possible to create UNPs that vary from nearly perfect fluids (Euler limit) to highly kinetic plasmas. We have examined plasmas in three geometries: a solid rod, a hollow rod and a gapped slab; we have studied basic properties of these plasmas, including the spatial Coulomb coupling parameter. By varying the initial conditions, we find that we can design experimental plasmas that would allow the exploration of a wide range of phenomena, including shock and blast waves, stopping power, two-stream instabilities and much more. Using an evaporative cooling geometry, our results suggest that much larger Coulomb couplings can be achieved, possibly in excess of ten.

References

[1] Vikram S Dharodi and Michael S Murillo. Sculpted ultracold neutral plasmas. *Physical Review E*, 101(2):023207, 2020.
