

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

84da148293bcb860ace1a4c83542b39833f20bb3f87f6584afdc1ae4db825120

To view the reconstructed contents, please SCROLL DOWN to next page.

# Seminar

---

---

## Institute for Plasma Research

---

---

**Title :** Role of nuclear dissipation in heavy ion fusion-fission reactions

**Speaker:** Dr. Niraj Kumar Rai  
Banaras Hindu University, Varanasi

**Date :** 30th December 2020 (Wednesday)

**Time :** 03.30 PM

**Venue :** Online - Join the talk:

[https://meet.ipr.res.in/Dr.NirajKumar\\_PDFTalk](https://meet.ipr.res.in/Dr.NirajKumar_PDFTalk)

### Abstract :

The fusion-fission process of the excited compound nucleus can be understood by the evaporation of particles such as alpha, proton, neutron, gamma, and fission fragments. In the collision between two heavy nuclei, quasi-fission as well as fusion-fission has a considerable contribution in the process. The measurement of the pre-scission multiplicity of light charged particles [1], the neutrons [2–3], and GDR  $\gamma$  rays [4], evaporation residue cross sections [5], and the mass and angular distribution of the fission fragments [6] are well-established tools to understand the heavy-ion induced fusion-fission process. Nuclear dissipation plays an important role in the nuclear physics research area. The presence of nuclear dissipation in the fusion-fission process is observed but it was not mentioned that how it depends on the entrance channel parameters. Therefore, it was studied about the effect of entrance channel parameters like projectile energy, mass asymmetry, and Coulomb factor ZPZT [7]. The pre- and post-scission neutron multiplicities were measured for the reactions  $180 + 186, 184W$  at different excitation energies populating the compound nucleus  $204, 202Pb$ , using the National Array of Neutron Detectors (NAND) facility at IUAC, New Delhi, India [8]. Here, it was investigated about the entrance channel effect on the nuclear dissipation involved in the heavy ion fusion-fission dynamics. Specifically, the role of the entrance channel parameters on the nuclear dissipation was studied.

### References:

- [1] J. P. Lestone, Phys. Rev. Lett. 70, 2245 (1993).
  - [2] K. Ramachandran et al., Phys. Rev. C 73, 064609 (2006).
  - [3] J. Cabrera et al., Phys. Rev. C 68, 034613 (2003).
  - [4] I. Dio'szegi et al., Phys. Rev. C 61, 024613 (2000).
  - [5] B. B. Back et al., Phys. Rev. C 60, 044602 (1999).
  - [6] A. Saxena et al., Phys. Rev. C 49, 932 (1994).
  - [7] N. K. Rai et al., Phys. Rev. C 98, 024626 (2018).
  - [8] N. K. Rai et al., Phys. Rev. C 100, 014614 (2019).
-