

This file has been cleaned of potential threats.

To view the reconstructed contents, please SCROLL DOWN to next page.

# Seminar

---

---

## Institute for Plasma Research

---

---

**Title:** Control of Edge and Scrape-off Layer Tokamak Plasma Turbulence  
**Speaker:** Mr. Vijay Shankar  
Institute for Plasma Research, Gandhinagar  
**Date:** 12<sup>th</sup> March 2024 (Tuesday)  
**Time:** 11:00 AM  
**Venue:** Seminar Hall, IPR  
Join the Seminar online: <https://meet.google.com/gvq-yzfr-kjx>

### Abstract

The plasma in the edge and scrape-off layer (SOL) regions of a tokamak is highly turbulent mainly due to the excitation of the interchange instability. This instability occurs as the directions of centrifugal force due to the rapid toroidal plasma motion and the density/pressure gradient are in opposing directions. The plasma turbulence gives rise to some coherent nonlinear density structures called “blobs” and these blobs are responsible for the anomalous transport of the plasma. Therefore, controlling anomalous transport and these instabilities is necessary for the stable/safe operation of tokamaks. Various processes have been used to control these instabilities in the past. In this work, we investigate the control of turbulence from the perspective of the blob birth generation mechanisms, and by applying an external bias (electrode biasing) in the edge region.

The formation of a blob in the edge and SOL regions of a tokamak plasma normally takes place when a radially elongated density streamer structure breaks due to differential stretching in the radial and poloidal directions. Such a phenomenon has been extensively studied in the past in the edge-to-SOL transition region using 2D (two-dimensional) model equations. In this work, we have examined the blob formation mechanism in the SOL and far SOL regions of a tokamak plasma using a 3D (three-dimensional) model theoretically as well as numerically. The blob formation mechanism is found to be quite different from those in the edge-to-SOL transition region where the poloidal gradient of the poloidal electric field (PGPEF) is playing an important role which is not considered in the earlier works. We have derived a heuristic condition for such a plasma blob formation theoretically. Blob formation takes place when the amount of shear exceeds the non-linear growth rate ( $\gamma_n$ ) of the basic interchange instability within the radially elongated streamer region. From 3D simulational data, it has been observed that the cross-correlation between PGPEF and  $\gamma_n$  is higher than the cross-correlation between other shears and  $\gamma_n$ . These correlations indicate that  $\gamma_n$  and these shears are not independent, they may be connected by a blob formation. In this work, the blob formation has been investigated in terms of a criterion using a large number of blob formations.

Electrode biasing in the Tokamak boundary region is an important topic as it can control particle recycling, exhausts, confinement time, and heat loads on the limiter/divertor plates. Here, in this work, we have investigated some of the above effects theoretically and numerically. A linear analysis of a set of model fluid equations shows that biasing stabilizes the small  $k_y$  modes. Positive biasing is found to lead to a larger increment in plasma density and temperature as compared to negative biasing. Nonlinearly, it is further observed that cross-correlation between density and poloidal electric field at different radial posi-

tions decreases for positive biasing, and in the case of negative biasing, it is almost similar to that of no (w/o) biasing. Plasma density and poloidal electric field fluctuations have been investigated which show that the density fluctuations increase (decrease) for positive (negative) biasing. The concomitant modifications in the power spectra are also analyzed. The positive bias shifts the peak of the  $k_y$ -spectra towards a lower  $k_y$  value in the edge region. The heat and particle fluxes in the edge region are seen to increase with the radial electric field shear in the region where a flow reversal takes place. Elsewhere, the fluxes decrease with an increase in the shear. The heat and particle load per unit time on the limiter/divertor material plates decrease with the biasing voltages in comparison to those w/o biasing, but the load intensities increase. It is found that the blob fraction decreases with the radial electric field shear but at lower radial positions around the biasing region it increases from -16 volts to +64 volts.

---